

South Island Notifiable Disease Rates By DHB*

Rates (per 100,000 population) for the 12 months to July 2019

[EpiSurv extract N010 extracted 1200 hrs on 6 August 2019]

	Nels Marl	West Coast	Canterbury	Sth Canty	Southern	NZ
Brucellosis	0.0	0.0	0.2	0.0	0.0	0.0
Campylobacteriosis	151.4	138.0	140.8	255.4	180.6	136.9
Chikungunya fever	0.0	0.0	0.5	0.0	0.3	0.2
Cryptosporidiosis	19.3	15.3	15.8	45.1	33.6	21.7
Cysticercosis	0.0	0.0	0.0	0.0	0.0	0.0
Dengue fever	4.6	0.0	2.3	5.0	2.4	3.6
Diphtheria	0.0	0.0	0.2	0.0	0.0	0.0
Gastroenteritis - unknown cause	2.7	3.1	5.1	5.0	1.8	3.7
Gastroenteritis / foodborne intoxication	2.7	3.1	1.2	1.7	0.6	4.7
Giardiasis	47.8	9.2	29.5	40.1	31.2	33.3
Haemophilus influenzae type b	0.0	0.0	0.0	0.0	0.0	0.1
Hepatitis A	0.7	0.0	0.9	0.0	0.3	1.4
Hepatitis B	0.7	0.0	0.4	0.0	0.0	0.8
Hepatitis C	2.7	6.1	1.4	0.0	2.1	0.8
Hepatitis NOS	0.7	0.0	0.0	0.0	0.0	0.2
Hydatid disease	0.0	0.0	0.0	1.7	0.0	0.0
Invasive pneumococcal disease	12.0	9.2	7.5	5.0	10.0	10.6
Legionellosis	1.3	3.1	5.7	0.0	3.9	2.9
Leprosy	0.0	0.0	0.0	0.0	0.3	0.1
Leptospirosis	3.3	12.3	0.5	1.7	1.5	1.9
Listeriosis	0.0	0.0	0.7	1.7	0.3	0.5
Listeriosis - perinatal	0.0	0.0	0.4	0.0	0.0	0.1
Malaria	1.3	0.0	0.7	0.0	0.3	0.7
Measles	0.7	0.0	7.3	0.0	1.2	9.1
Meningococcal disease	2.0	0.0	2.0	1.7	1.5	3.0
Mumps	2.0	0.0	0.5	0.0	1.2	2.6
Paratyphoid fever	0.0	3.1	0.4	0.0	0.0	0.3
Pertussis	112.9	199.4	16.3	0.0	17.6	40.3
Rheumatic fever - initial attack	0.7	0.0	0.2	0.0	0.6	3.2
Rheumatic fever - recurrent attack	0.0	0.0	0.0	0.0	0.0	0.3
Rickettsial disease	0.0	0.0	0.0	0.0	0.0	0.1
Ross River virus infection	0.0	0.0	0.0	0.0	0.0	0.1
Salmonellosis	27.2	52.1	25.9	30.1	42.4	24.1
Shigellosis	7.3	0.0	2.3	0.0	1.8	4.6
Taeniasis	0.0	0.0	0.0	0.0	0.0	0.2
Toxic shellfish poisoning	0.7	0.0	0.0	0.0	0.0	0.0
Tuberculosis disease - new case	2.7	9.2	5.3	6.7	2.7	6.8
Typhoid fever	0.7	0.0	0.9	0.0	0.6	1.1
VTEC/STEC infection	33.9	0.0	11.4	11.7	60.0	21.9
Yersiniosis	27.9	6.1	24.5	8.3	21.5	21.6

*Zika notifications are only being reported nationally