

Te Waipounamu Notifiable Disease Rates by District

Rates (per 100,000 population) for the 12 months to October 2024

[EpiSurv extract N010 extracted 0750 hrs on 13 November 2024]

| | Nelson Marlborough | Te Tai o Poutini West Coast | Waitaha Canterbury | South Canterbury | Southern | New Zealand |
|--|-----------------------|--------------------------------|-----------------------|---------------------|----------|-------------|
| Botulism | 0 | 0 | 0.2 | 0 | 0 | 0 |
| Campylobacteriosis | 102.3 | 139.8 | 108.6 | 192.1 | 138.9 | 109.4 |
| Chikungunya fever | 0 | 0 | 0.2 | 0 | 0.3 | 0.1 |
| Cholera | 0 | 0 | 0.2 | 0 | 0 | 0.1 |
| Cronobacter species | 0 | 0 | 0 | 0 | 0 | 0 |
| Cryptosporidiosis | 9 | 18.2 | 37.2 | 54 | 24.4 | 24.4 |
| Decompression sickness | 0 | 0 | 0.2 | 0 | 0 | 0 |
| Dengue fever | 1.8 | 3 | 2.2 | 0 | 1.4 | 2.2 |
| Diphtheria | 0 | 0 | 0 | 1.6 | 0 | 0 |
| Gastroenteritis - unknown cause | 3 | 12.2 | 10.1 | 1.6 | 8.3 | 3.8 |
| Gastroenteritis / foodborne intoxication | 0 | 15.2 | 4 | 1.6 | 0.3 | 4.3 |
| Giardiasis | 6 | 12.2 | 9.8 | 7.9 | 11.6 | 16.6 |
| Haemophilus influenzae type b | 0 | 0 | 0.2 | 0 | 0 | 0 |
| Hepatitis A | 0.6 | 0 | 2.3 | 0 | 0.6 | 1.1 |
| Hepatitis B | 0.6 | 0 | 0.2 | 0 | 0.3 | 0.3 |
| Hepatitis C | 1.2 | 3 | 0.2 | 1.6 | 1.1 | 0.6 |
| Hepatitis NOS | 0 | 0 | 0.2 | 0 | 0 | 0.1 |
| Hydatid disease | 0 | 0 | 0 | 0 | 0 | 0.1 |
| Invasive group A streptococcal infection | 1.2 | 3 | 0.7 | 1.6 | 1.9 | 0.9 |
| Invasive pneumococcal disease | 11.4 | 21.3 | 10.5 | 6.3 | 12.2 | 14 |
| Latent tuberculosis infection | 0.6 | 3 | 7.5 | 6.3 | 0 | 2.3 |
| Legionellosis | 3 | 3 | 8 | 4.8 | 5.5 | 4.2 |
| Leprosy | 0 | 0 | 0.2 | 0 | 0.3 | 0.1 |
| Leptospirosis | 4.2 | 15.2 | 0.2 | 1.6 | 1.1 | 2.1 |
| Listeriosis | 0 | 0 | 0.7 | 1.6 | 1.1 | 0.5 |
| Listeriosis - perinatal | 0 | 0 | 0.2 | 0 | 0 | 0.1 |
| Malaria | 0.6 | 0 | 0 | 0 | 0.8 | 0.6 |
| Measles | 0 | 0 | 0 | 0 | 0 | 0 |
| Meningococcal disease | 0 | 3 | 0.3 | 0 | 0.6 | 0.8 |
| Mumps | 0 | 0 | 0.8 | 1.6 | 0.8 | 0.5 |
| Murine Typhus | 0 | 0 | 0 | 0 | 0 | 0 |
| Paralytic shellfish poisoning | 0 | 0 | 0 | 0 | 0 | 0 |
| Paratyphoid fever | 0 | 3 | 0.5 | 0 | 0 | 0.3 |
| Pertussis | 6 | 3 | 19.8 | 9.5 | 8 | 15.2 |
| Rheumatic fever - initial attack | 0 | 0 | 1.5 | 1.6 | 1.1 | 3.6 |
| Rheumatic fever - recurrent attack | 0 | 0 | 0.2 | 0 | 0 | 0.4 |
| Rickettsial disease | 0 | 0 | 0 | 0 | 0 | 0.1 |
| Ross River virus infection | 0 | 0 | 0 | 0 | 0.3 | 0 |
| Salmonellosis | 8.4 | 3 | 18.9 | 31.7 | 20 | 15 |
| Shigellosis | 2.4 | 9.1 | 2 | 6.3 | 1.1 | 3.3 |
| Taeniasis | 0 | 0 | 0.2 | 1.6 | 0 | 0 |
| Tetanus | 0 | 0 | 0 | 0 | 0 | 0 |
| Toxic shellfish poisoning | 0 | 0 | 0 | 0 | 0 | 0 |
| Tuberculosis disease - new case | 1.2 | 0 | 6.8 | 6.3 | 3.1 | 7 |
| Typhoid fever | 0.6 | 6.1 | 0.8 | 1.6 | 0.6 | 1.1 |
| VTEC/STEC infection | 25.7 | 27.4 | 25.7 | 76.2 | 42.2 | 21.8 |
| Yersiniosis | 33.5 | 15.2 | 24.3 | 31.7 | 30.5 | 22 |

COVID-19 information is being reported by the Ministry of Health nationally.

Zika notifications are reported nationally and mpox notifications by region (i.e. Te Waipounamu/South Island).